

ON TAMENESS AND GROWTH CONDITIONS

JÖRG WINKELMANN

ABSTRACT. We study discrete subsets of $\mathbb{C}^d$, relating “tameness” with growth conditions.

1. RESULTS

A discrete subset D in $\mathbb{C}^n$ ($n \geq 2$) is called “tame” if there exists a holomorphic automorphism ϕ of $\mathbb{C}^n$ such that $\phi(D) = \mathbb{Z} \times \{0\}^{n-1}$ (see [3]). If there exists a linear projection π of $\mathbb{C}^n$ onto some $\mathbb{C}^k$ ($0 < k < n$) for which the image $\pi(D)$ is discrete, then D is tame ([3]). If D is a discrete subgroup (e.g. a lattice) of the additive group $(\mathbb{C}^n, +)$, then D must be tame ([1], lemma 4.4 in combination with corollary 2.6). On the other hand there do exist discrete subsets which are not tame (see [3], theorem 3.9).

Here we will investigate how “tameness” is related to growth conditions for D .

Slow growth implies tameness as we will see. On the other hand, rapid growth can not imply non-tameness, since every discrete subset of $\mathbb{C}^{n-1}$ is tame regarded as subset of $\mathbb{C}^n = \mathbb{C}^{n-1} \times \mathbb{C}$.

The key method is to show that sufficiently slow growth implies that a generic linear projection will have discrete image for D .

The main result is:

Theorem 1. *Let n be a natural number and let v_k be a sequence of elements in $V = \mathbb{C}^n$.*

Assume that

$$\sum_k \frac{1}{\|v_k\|^{2n-2}} < \infty$$

Then $D = \{v_k : k \in \mathbb{N}\}$ is tame, i.e., there exists a biholomorphic map $\phi : \mathbb{C}^n \rightarrow \mathbb{C}^n$ such that

$$\phi(D) = \mathbb{Z} \times \{0\}^{n-1}.$$

This growth condition is fulfilled for discrete subgroups of rank at most $2n - 3$, implying the following well-known fact:

Corollary 1. *Let Γ be a discrete subgroup of $\mathbb{Z}$ -rank at most $2n - 3$ of the additive group $(\mathbb{C}^n, +)$.*

Then Γ is a tame discrete subset of $\mathbb{C}^n$.

While this is well-known (even with no condition on the $\mathbb{Z}$ -rank of Γ), our approach yields the additional information that these discrete subsets remain tame after a small deformation:

Corollary 2. *Let Γ be a discrete subgroup of $\mathbb{Z}$ -rank at most $2n - 3$ of the additive group $(\mathbb{C}^n, +)$, $0 < \lambda < 1$ and $K > 0$. Let D be a subset of $\mathbb{C}^n$ for which there exists a bijective map $\zeta : \Gamma \rightarrow D$ with*

$$\|\zeta(v) - v\| \leq \lambda\|v\| + K$$

for all $v \in \Gamma$.

Then D is a tame discrete subset of $\mathbb{C}^n$.

This confirms the idea that tame sets should be stable under deformation. Similarly one would hope that the category of non-tame sets is also stable under deformation. Here, however, one has to be careful not to be too optimistic, because in fact the following is true:

Proposition. *For every non-tame discrete subset $D \subset \mathbb{C}^n$ ($n > 1$) there is a tame discrete subset D' and a bijection $\alpha : D \rightarrow D'$ such that*

$$\|\alpha(v) - v\| \leq \frac{1}{\sqrt{2}}\|v\| \quad \forall v \in D$$

and

$$\|w - \alpha^{-1}(w)\| \leq \|w\| \quad \forall w \in D'.$$

In particular, if D is a tame discrete subset and $\zeta : D \rightarrow \mathbb{C}^n$ is a bijective map with $\|\zeta(v) - v\| \leq \|v\|$ for all $v \in D$, it is possible that $\zeta(D)$ is not tame.

Still, one might hope for a positive answer to the following question:

Question. *Let $n \in \mathbb{N}$, $n \geq 2$, let $1 > \lambda > 0$, $K > 0$, let D be a tame discrete subset of $\mathbb{C}^n$ and let $\zeta : D \rightarrow \mathbb{C}^n$ be a map such that*

$$\|\zeta(v) - v\| \leq \lambda\|v\| + K$$

for all $v \in D$. Does this imply that $\zeta(D)$ is a tame discrete subset of $\mathbb{C}^n$?

Technically, the following is the key point for the proof of our main result (theorem 1):

Theorem 2. *Let $n > d > 0$. Let V be a complex vector space of dimension n and let v_k be a sequence of elements in V .*

Assume that

$$\sum_k \frac{1}{||v_k||^{2d}} < \infty$$

Then there exists a complex linear map $\pi : V \rightarrow \mathbb{C}^d$ such that the set of all $\pi(v_k)$ is discrete in $\mathbb{C}^d$.

In a similar way one can prove such a result for real vector spaces:

Theorem 3. *Let $n > d > 0$. Let V be a real vector space of dimension n and let v_k be a sequence of elements in V .*

Assume that

$$\sum_k \frac{1}{||v_k||^d} < \infty$$

Then there exists a real linear map $\pi : V \rightarrow \mathbb{R}^d$ such that the set of all $\pi(v_k)$ is discrete in $\mathbb{R}^d$.

For the proof of the existence of a linear projection π with $\pi(D)$ discrete we proceed by regarding randomly chosen linear projections and verifying that the image of D under a random projection has discrete image with probability 1 if the above stated series converges.

2. PROOFS

First we deduce an auxiliary lemma.

Lemma 1. *Let $k, m > 0$, $n = k + m$ and let S denote the unit sphere in $\mathbb{R}^n = \mathbb{R}^k \oplus \mathbb{R}^m$. Furthermore let*

$$M_\epsilon = \{(v, w) \in \mathbb{R}^k \times \mathbb{R}^m : ||v|| \leq \epsilon, (v, w) \in S\}.$$

Then there are constants $\delta > 0$, $C_1 > C_2 > 0$ such that for all $\epsilon < \delta$ we have

$$C_1 \epsilon^k \geq \lambda(M_\epsilon) \geq C_2 \epsilon^k$$

where λ denotes the rotationally invariant probability measure on S .

Proof. For each $\epsilon \in]0, 1[$ there is a bijection

$$\phi_\epsilon : B \times S' \rightarrow M_\epsilon$$

where

$$B = \{v \in \mathbb{R}^k : ||v|| \leq 1\}, \quad S' = \{w \in \mathbb{R}^m : ||w|| = 1\}$$

and

$$\phi_\epsilon(v, w) = \left(\epsilon v; \sqrt{1 - ||\epsilon v||^2} w \right).$$

The functional determinant for ϕ_ϵ equals

$$\epsilon^k \left(\sqrt{1 - ||\epsilon v||^2} \right)^m.$$

It follows that

$$\epsilon^k \left(\sqrt{1 - \epsilon^2} \right)^m \text{volume}(S' \times B) \leq \text{volume}(M_\epsilon) \leq \epsilon^k \text{volume}(S' \times B),$$

which in turn implies

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-k} \frac{\text{volume}(M_\epsilon)}{\text{volume}(S' \times B)} = 1.$$

Hence the assertion. $\square$

Lemma 2. *Let Γ be a discrete subgroup of $\mathbb{Z}$ -rank d in $V = \mathbb{R}^n$.*

Then

$$\sum_{\gamma \in \Gamma} \|\gamma\|^{-d-\epsilon} < \infty$$

for all $\epsilon > 0$.

Proof. Since all norms on a finite-dimensional vector space are equivalent, there is no loss in generality if we assume that the norm is the maximum norm and $\Gamma = \mathbb{Z}^d \times \{0\}^{n-d}$. Then the assertion is an easy consequence of the fact that $\sum_{n \in \mathbb{N}} n^{-s} < \infty$ if and only if $s > 1$. $\square$

Now we proceed with the proof of theorem 2:

Proof. We fix a surjective linear map $L : V \rightarrow W = \mathbb{C}^d$. Let K denote $U(n)$ (the group of unitary complex linear transformations of V). For each $g \in K$ we define a linear map $\pi_g : V \rightarrow W$ as follows:

$$\pi_g : v \mapsto L(g \cdot v).$$

For $k \in \mathbb{N}$ and $r \in \mathbb{R}^+$ define

$$S_{k,r} = \{g \in K : \|\pi_g(v_k)\| \leq r\},$$

$$M_{N,r} = \{g \in K : \#\{k \in \mathbb{N} : g \in S_{k,r}\} \geq N\}$$

and

$$M_r = \cap_N M_{N,r}.$$

Now for each $g \in K$ the set $\{\pi_g(v_k) : k \in \mathbb{N}\}$ is discrete unless there is a number $r > 0$ such that infinitely many distinct image points are contained in a ball of radius r . By the definition of the sets M_r it follows that $\{\pi_g(v_k) : k \in \mathbb{N}\}$ is discrete unless $g \in M = \cup M_r$.

Let us now assume that there is no linear map $L' : V \rightarrow W$ with $L'(D)$ discrete. Then $K = M$. In particular $\mu(M) > 0$, where μ denotes the Haar measure on the compact topological group K . Since the sets M_r are increasing in r , we have

$$M = \cup_{r \in \mathbb{R}^+} M_r = \cup_{r \in \mathbb{N}} M_r$$

and may thus deduce that $\mu(M_r) > 0$ for some number r . Fix such a number $r > 0$ and define $c = \mu(M_r) > 0$. Then $\mu(M_{N,r}) \geq c$ for all N , since $M_r = \cap M_{N,r}$. However, for fixed N and r we have

$$N\mu(M_{N,r}) \leq \sum_k \mu(S_{k,r}).$$

Hence

$$\sum_{k \in \mathbb{N}} \mu(S_{k,r}) \geq N\mu(M_{N,r}) \geq Nc$$

for all $N \in \mathbb{N}$. Since $c > 0$, it follows that $\sum_k \mu(S_{k,r}) = +\infty$.

Let us now embed $\mathbb{C}^d$ into $\mathbb{C}^n$ as the orthogonal complement of $\ker L$. In this way we may assume that L is simply the map which projects a vector onto its first d coordinates, i.e.,

$$L(w_1, \dots, w_n) = (w_1, \dots, w_d; 0, \dots, 0).$$

Now $g \in S_{k,r}$ is equivalent to the condition that $g(v_k)$ is a real multiple of an element in M_ϵ where M_ϵ is defined as in lemma 1 with $\epsilon = r/||v_k||$. Using lemma 1 we may deduce that $\sum_k \mu(S_{k,r})$ converges if and only if $\sum_k ||v_k||^{-2d}$ converges. $\square$

Proof of theorem 1. The growth condition allows us to employ theorem 2 in order to deduce that there is a linear projection onto a space of complex dimension $d - 1$ which maps D onto a discrete image. By the results of Rosay and Rudin it follows that D is tame. $\square$

Proof of the proposition. We fix a decomposition $\mathbb{C}^n = \mathbb{C} \times \mathbb{C}^{n-1}$ and write D as the union of all $(a_k, b_k) \in \mathbb{C} \times \mathbb{C}^{n-1}$ ($k \in \mathbb{N}$). We define

$$\alpha(a_k, b_k) = \begin{cases} (a_k, 0) & \text{if } ||a_k|| > ||b_k|| \\ (0, b_k) & \text{if } ||a_k|| \leq ||b_k|| \end{cases}$$

Then $D' = \alpha(D)$ is tame because each of the projections to one of the two factors $\mathbb{C}$ and $\mathbb{C}^{n-1}$ maps D' onto a discrete subset.

The other assertions follow from the triangle inequality. $\square$

The proof of thm. 3 works in the same way as the proof of thm. 2, simply using the group of all orthogonal transformations instead of the group of unitary transformations.

REFERENCES

- [1] Buzzard, G.: Tame sets, dominating maps and complex tori. *Trans. A.M.S.* **355**, 2557-2568 (2002)
- [2] Buzzard, G.; Forstneric, F.: An interpolation theorem for holomorphic automorphisms of $\mathbb{C}^n$. *J. Geom. Anal.* **10**, 101-108 (2000)

- [3] Rosay, J.P.; Rudin, W.: Holomorphic maps from $\mathbb{C}^n$ to $\mathbb{C}^n$. *Trans. A.M.S.* **310**, 47–86 (1988)

JÖRG WINKELMANN, MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH,
UNIVERSITÄTSSTRASSE 30, D-95447 BAYREUTH, GERMANY,

E-mail address: `jwinkel@member.ams.org`

Webpage: `http://btm8x5.mat.uni-bayreuth.de/~winkelmann/`